# **Atlantic Whitefish**


Coregonus huntsmani





## **Species Description**

The Atlantic Whitefish has an elongated body, a fleshy adipose fin, and a deeply forked caudal fin (tail). It has a dark green to dark blue back, slightly lighter sides and a silvery white belly. It has dusky dorsal and caudal fins and light pelvic and anal fins. It is typically 20-25 cm long.



- Terminal mouth (faces forward at end of snout)
- More than 90 large lateral line scales
- Upper and lower jaws fairly equal in length
- No spots or upper body markings

Only one small population in the world, and it is in the Petite Rivière watershed in NS. Considered extirpated from the Tusket River, NS.



Until recently, Atlantic Whitefish were confined to three small seminatural, inter-connected lakes (Hebb, Milipsigate and Minamkeak Lakes) as a result of a dam located at the base of the lakes. In 2012, a fishway was constructed at the Hebb Dam in an attempt to restore conditions that allow access to the ocean. In freshwater, this species is found throughout the water column, with adults generally found in cooler, deeper water and juveniles in shallower water.





Atlantic Whitefish are an anadromous (sea run) fish by nature, which means they are meant to spawn in freshwater then return to the ocean to feed, grow and mature.

### **Interesting Points**

- The most primitive of all North American whitefishes (oldest evolutionary link).
- Found nowhere else in the world!
- Also known as the Acadian, Sault or Round Whitefish.
- Old stories say that Tusket River Atlantic Whitefish were so plentiful they were taken by the truckload and used as fertilizer or lobster bait.
- Prior to their decline, larger bodied sea-run individuals could reach lengths of up to 50 cm.
- Atlantic Whitefish raised in captivity were introduced into a small lake in Dartmouth (Anderson Lake), however it is currently not known whether they have established a population.

# **Similar Species**

#### Lake Whitefish:

Common, smaller and fewer than 90 lateral scales (on side from head to tail); subterminal mouth (faces downwards).





Newly constructed fishway



### Threats to Survival

- Fish passage barriers impede access to the sea (several dams still exist in the Petite Rivière watershed).
- Unauthorized introduction of invasive alien species (Chain Pickerel, Smallmouth Bass).
- Increased acidification of lake water from acid rain and landbased activities (forestry, agricultural, residential).
- Accidental catch by recreational anglers.



Never move fish from one lake into another

## **How You Can Help**

Prevent the introduction of invasive species; it is illegal to move fish from one body of water to another. Fishing Atlantic Whitefish is prohibited. Learn to identify this species and if accidentally caught, notify Bluenose Coastal Action Foundation (BCAF) to report a sighting. Protect fish habitat on your land by leaving a buffer of trees along the waters edge, and avoid the use of chemicals and fertilizers.



Monitoring migration downstream of the fishway

#### Contacts, Information, Sighting Reports & Stewardship Opportunities

Contact: DFO Species at Risk I-866-891-0771or xmarsara@mar.dfo-mpo.gc.ca, and NS Department of Agriculture and Fisheries inlandfish@gov.ns.ca

Info: www.sararegistry.gc.ca, www.dfo-mpo.gc.ca/species-especes, www.novascotia.ca/fish

Sighting Reports: 1-866-727-3447 or www.speciesatrisk.ca/sightings Stewardship: BCAF (902) 634-9977 www.coastalaction.org